

PrimeGene Technical Data Sheet

Catalog Number:	105-44
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 34.6 kDa, a disulfide-linked homodimer consisting of two 152 amino acid polypeptide chains.
Quantity:	5µg/25µg/1000µg
AA Sequence:	LPAVPPQQWA LSAGNGSSEV EVVPFQEVWG RSYCRALERL VDVVSEYPSV VEHMFSPSCV SLLRCTGCCG DENLHCVPVE TANVTMQLLK IRSGDRPSYV ELTFSQHVRV ECRPLREKMK PERRRPKGRG KRRREKQRPT DCHLCGDAVP RR
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biologically active as determined by its ability to chemoattract human monocytes using a concentration range of 5.0-50 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4, with 0.02 % Tween-20.
Endotoxin:	Less than 0.1 EU/µg of rHuPIGF-2 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Placenta Growth Factor-2

Placenta growth factor (PIGF) is a member of the vascular endothelial growth factor (VEGF) family of growth factors. PIGF and VEGF share primary structural as well as limited amino acid sequence homology with the A and B chains of PDGF. All eight cysteine residues involved in intra-and inter-chain disulfides are conserved among these growth factors. As a result of alternative splicing, three PIGF RNAs encoding monomeric human PIGF-1, PIGF-2 and PIGF-3 isoform precursors containing 149, 179 and 219 amino acid residues, respectively, have been described. Human PIGF-2 shares 65 % amino acid identity with murine PIGF-2.