

Prime Gene Recombinant Human Ciliary Neurotrophic Factor (rHuCNTF)

PrimeGene Technical DataSheet

Catalog Number: 107-08G

Source: Escherichia coli

Molecular Weight: Approximately 22.9 kDa, a single non-glycosylated polypeptide chain containing 200 amino acids.

Size: 5ug/100ug/500ug/1mg

AA Sequence: MAFTEHSPLT PHRRDLCSRS IWLARKIRSD LTALTESYVK HOGLNKNINL DSADGMPVAS

> TDQWSELTEA ERLQENLQAY RTFHVLLARL LEDQQVHFTP TEGDFHQAIH TLLLQVAAFA YQIEELMILL EYKIPRNEAD GMPINVGDGG LFEKKLWGLK VLQELSQWTV RSIHDLRFIS

SHOTGIPARG SHYIANNKKM

Purity: \geq 97 % by SDS-PAGE analysis. **Biological Activity:** Bioassay data are not available.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a 0.2 µm filtered solution in 1 × PBS, pH 7.0, with 5 % Trehalose, 0.02 % Tween-

20, 0.02 % Tween-80, 1 mM EDTA.

Endotoxin: Less than 0.1 EU/µg of rHuCNTF as determined by LAL method.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the

> bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and

stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.

Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature

recommended below.

Stability & Storage: Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70 °C as supplied.

1 month, 2 to 8 °C under sterile conditions after reconstitution.

3 months, -20 to -70 °C under sterile conditions after reconstitution.

Usage: This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further

evaluation purposes. NOT FOR HUMAN USE.

Human Ciliary Neurotrophic Factor

Ciliary neurotrophic factor (CNTF) is a polypeptide hormone whose actions appear to be restricted to the nervous system where it promotes neurotransmitter synthesis and neurite outgrowth in certain neuronal populations. It was initially identified as a trophic factor for embryonic chick ciliary parasympathetic neurons in culture. The protein is also a potent survival factor for neurons and oligodendrocytes and may be relevant in reducing tissue destruction during inflammatory attacks. In addition, CNTF is useful for treatment of motor neuron disease, and it could reduce food intake without causing hunger or stress. CNTF is structurally related to IL-6, IL-11, LIF and OSM. All of these four helix bundle cytokines share gp130 as a signal-transducing subunit in their receptor complexes. Recombinant human CNTF containing 200 amino acids and it shares 82 % and 83 % a.a. sequence identity with mouse and rat CNTF.

Shanghai PrimeGene Bio-Tech Co., Ltd.

Rev. 03/27/2025 V.2 Email: info.pg@bio-techne.com Website: www.primegene.com.cn Website: www.primegene.com

Tel: +86 21 52293200