

**Recombinant Bifunctional ligase/repressor
BirA, His
(rBirA, His)
PrimeGene Technical Data Sheet**

Catalog Number:	4A8-01H
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 36.4 kDa, a single non-glycosylated polypeptide chain containing 329 amino acids, with 6 × His at the C-terminus.
Quantity:	5µg/100µg/500µg
AA Sequence:	MKDNTVPLKL IALLANGEFH SGEQLGETLG MSRAAINKHI QTLRDWGVVDV FTVPGKGYSL PEPIQLLNAK QILGQLDGGG VAVLPVIDST NQYLLDRIGE LKSGDACIAE YQQAGRGRRG RKWFSPFGAN LYLSMFWRL E QGPAAIGLS LVIGIVMAEV LRKLGADKVR VKWPNDLYLQ DRKLAGILVE LTGKTGDA AQ IVIGAGINMA MRRVEESVVN QGWITLQEAG INLDRNTLAA MLIRELRAAL ELFEQEGLAP YLSRWEKLDN FINRPVKLII GDKEIFGISR GIDKQGALL EQDGIKPWM GGEISLRS AE KLEHHHHHHH
Purity:	> 95 % by SDS-PAGE analyses.
Biological Activity:	Measured by its ability to generate pyrophosphate from the biotinylation reaction. The pyrophosphate is subsequently hydrolyzed using Recombinant Yeast Inorganic Pyrophosphatase/PPA1 (ryPPA1). The specific activity is > 10.0 pmol/min/µg, as measured under the described conditions.
Physical Appearance:	Sterile colorless liquid.
Formulation:	Supplied as a 0.2 µm filtered solution in 50 mM Tris-HCl, pH 8.0, 50 mM NaCl, 1 mM DTT, 10 % glycerol.
Endotoxin:	Less than 0.1 EU/µg of rBirA, His as determined by LAL method.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 6 months from date of receipt, -20 to -70 °C as supplied.● 3 months, -20 to -70 °C under sterile conditions after opening.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Bifunctional ligase/repressor BirA

BirA, the biotin-protein ligase (BPL) of *Escherichia coli*, is also known as biotin operon repressor, biotin-[acetyl-CoA-carboxylase] ligase, and biotin-[acetyl-CoA-carboxylase] synthetase. BirA, a member of the group II biotin-protein ligase family, contains an N-terminal helix-turn-helix DNA-binding domain, a catalytic core that catalyzes biotinyl 5' adenylate (bio-5'-AMP) synthesis, and a C-terminal domain that plays a role in DNA binding, dimerization, and catalytic function. BirA functions both as a DNA-binding protein that represses the biotin biosynthesis operon as well as an enzyme that synthesizes its own corepressor, bio-5'-AMP, an intermediate in biotinylation reactions. BirA biotinylates via the lysine side chain of biotin-accepting proteins/peptides, including natural substrate, carboxyl carrier protein (BCCP), and Avi Tag fusion proteins. Once biotinylated, (strept)avidin-biotin interactions can be utilized in a wide variety of applications of biochemistry and cell biology, including protein capture, immobilization, multimerizing, and bridging molecules.